

A NOTE ON SELF-EXTREMAL SETS IN $L_p(\Omega)$ SPACESby Viet Nguyen-Khac ¹ & Khiem Nguyen-Van ²

Hanoi Institute of Mathematics & Hanoi University of Education

1991 Mathematics Subject Classification. Primary 46B20, 46E30.

Abstract

We give a geometric characterization of self-extremal sets in $L_p(\Omega)$ spaces that partially extends our previous results to the case of L_p spaces.

In [5], [6] we introduced the notion of (self-) extremal sets of a Banach space $(X, \|\cdot\|)$. For a non-empty bounded subset A of X we denote by $r(A)$ the relative Chebyshev radius of A with respect to the closed convex hull $\overline{\text{co}}A$ of A , i.e. $r(A) := \inf_{y \in \overline{\text{co}}A} \sup_{x \in A} \|x - y\|$. The self-Jung constant of X is defined by $J_s(X) := \sup \{r(A) : A \subset X, \text{ with } \text{diam}(A) = 1\}$. If in this definition we replace $r(A)$ by the relative Chebyshev radius $r_X(A)$ of A with respect to the whole X , we get the Jung constant $J(X)$ of X . Recall that a bounded subset A of X consisting of at least two points is said to be extremal (resp. self-extremal) if $r_X(A) = J(X) d(A)$ (resp. $r(A) = J_s(X) d(A)$).

Let (Ω, μ) be a σ -finite measure space such that $L_p(\Omega)$ is infinite-dimensional. The Jung and self-Jung constants of $L_p(\Omega)$ ($1 \leq p < \infty$) were determined in [2], [4], [7], [8]:

$$J(L_p(\Omega)) = J_s(L_p(\Omega)) = \max \{2^{\frac{1}{p}-1}, 2^{-\frac{1}{p}}\}.$$

Theorem 1.1. *Let (Ω, μ) be a σ -finite measure space such that $L_p(\Omega)$ is infinite-dimensional ($1 < p < \infty$). If A is a self-extremal set in $L_p(\Omega)$, then $\alpha(A) = d(A)$.*

Here $\alpha(A) := \inf \{\varepsilon > 0 : A \text{ can be covered by finitely many sets of diameter } \leq \varepsilon\}$ - the Kuratowski measure of non-compactness of A .

Before proving our theorem we need the following results which for convenience we reformulate in the form of Lemmas 1.2 and 1.3 below.

¹Institute of Mathematics, 18 Hoang Quoc Viet, 10307 Hanoi, Vietnam

²Department of Mathematics & Informatics, Hanoi Univ. of Education, Cau Giay dist., Hanoi, Vietnam.

Lemma 1.2 ([2], Theorem 1.1). *Let X be a reflexive strictly convex Banach space and A a finite subset of X . Then there exists a subset $B \subset A$ such that*

- (i) $r(B) \geq r(A)$;
- (ii) $\|x - b\| = r(B)$ for every $x \in B$, where b is the relative Chebyshev center of B , i.e. $b \in \overline{\text{co}}B$ and $\sup_{x \in B} \|x - b\| = r(B)$.

Lemma 1.3 ([9], Theorem 15.1). *Let (Ω, μ) be a σ -finite measure space, $1 < p < \infty$, $x_1, \dots, x_n$ vectors in $L_p(\Omega)$ and $t_1, \dots, t_n$ non-negative numbers such that $\sum_{i=1}^n t_i = 1$. The following inequality holds*

$$2 \sum_{i=1}^n t_i \left\| x_i - \sum_{j=1}^n t_j x_j \right\|^\alpha \leq \sum_{i,j=1}^n t_i t_j \|x_i - x_j\|^\alpha,$$

where

$$\alpha = \begin{cases} q := \frac{p}{p-1} & \text{if } 1 < p \leq 2 \\ p & \text{if } p \geq 2 \end{cases} \quad (1)$$

Proof of Theorem 1.1. We may assume that A is closed convex and $r(A) = 1$. For each integer $n \geq 2$ we have

$$\bigcap_{x \in A} B(x, 1 - \frac{1}{n}) \cap A = \emptyset,$$

where $B(x, r)$ denotes the closed ball centered at x with radius r which is weakly compact since $L_p(\Omega)$ is reflexive. Hence there exist $x_{t_{n-1}+1}, x_{t_{n-1}+2}, \dots, x_{t_n}$ in A (with convention $t_1 = 0$) such that

$$\bigcap_{i=t_{n-1}+1}^{t_n} B(x_i, 1 - \frac{1}{n}) \cap A = \emptyset$$

Set $A_n := \{x_{t_{n-1}+1}, x_{t_{n-1}+2}, \dots, x_{t_n}\}$. By Lemma 1.2 there exists a subset $B_n = \{y_{s_{n-1}+1}, y_{s_{n-1}+2}, \dots, y_{s_n}\}$ of A_n satisfying properties (i)-(ii) of the lemma. Let us denote the relative Chebyshev center of B_n by c_n and let $r_n := r(B_n)$. By what we said above we have $r_n > 1 - \frac{1}{n}$ and $\|y_i - c_n\| = r_n$ for every $i \in I_n := \{s_{n-1}+1, s_{n-1}+2, \dots, s_n\}$. Since B_n is a finite set there exist non-negative numbers $\alpha_{s_{n-1}+1}, \alpha_{s_{n-1}+2}, \dots, \alpha_{s_n}$ with $\sum_{i \in I_n} \alpha_i = 1$ such that $c_n = \sum_{i \in I_n} \alpha_i y_i$. Applying Lemma 1.3 one gets

$$2r_n^\alpha = 2 \sum_{i \in I_n} \alpha_i \left\| y_i - \sum_{j \in I_n} \alpha_j y_j \right\|^\alpha \leq \sum_{i, j \in I_n} \alpha_i \alpha_j \|y_i - y_j\|^\alpha \quad (2)$$

where α as in (1).

Setting $B_\infty := \{y_{s_{n-1}+1}, y_{s_{n-1}+2}, \dots, y_{s_n}\}_{n=2}^\infty$ we claim that $\alpha(B_\infty) = d(A)$. Suppose on the contrary $\alpha(A_\infty) < d(A)$, so there exists $\varepsilon_0 \in (0, d(A))$ satisfying $\alpha(B_\infty) \leq d(A) - \varepsilon_0$, and subsets $D_1, D_2, \dots, D_m$ of $L_p(\Omega)$ with $d(D_i) \leq d(A) - \varepsilon_0$ for every $i = 1, 2, \dots, m$, such that $B_\infty \subset \bigcup_{i=1}^m D_i$. Then one can find at least one set among $D_1, D_2, \dots, D_m$, say D_1 with the property that there are infinitely many n satisfying

$$\sum_{i \in J_n} \alpha_i \geq \frac{1}{m} \quad (3)$$

where

$$J_n := \{i \in I_n : y_i \in D_1\}.$$

In view of (2) and taking into account $(d(A))^\alpha = \left(\frac{1}{J_s(L_p(\Omega))}\right)^\alpha = 2$ we have for all n satisfying (3)

$$\begin{aligned} 2r_n^\alpha &\leq \sum_{i, j \in I_n} \alpha_i \cdot \alpha_j \cdot \|y_i - y_j\|^\alpha \\ &\leq (d(A) - \varepsilon_0)^\alpha \cdot \left(\sum_{i, j \in J_n} \alpha_i \cdot \alpha_j \right) + (d(A))^\alpha \cdot \left(1 - \sum_{i, j \in J_n} \alpha_i \cdot \alpha_j \right) \\ &\leq 2 - \left[(d(A))^\alpha - (d(A) - \varepsilon_0)^\alpha \right] \cdot \frac{1}{m^2} \end{aligned} \quad (4)$$

On the other hand, obviously $1 - \frac{1}{n} < r_n \leq 1$, therefore $\lim_{n \rightarrow \infty} r_n = 1$. We get a contradiction with (4), since there are infinitely many n satisfying (3).

One concludes that $\alpha(B_\infty) = d(A)$, and hence $\alpha(A) = d(A)$.

The proof of Theorem 1.1 is complete.

As an immediate consequence one obtains an extension of Gulevich's result for $L_p(\Omega)$ spaces.

Corollary 1.4 (cf. [3]). *Let (Ω, μ) be a σ -finite measure space such that $L_p(\Omega)$ is infinite-dimensional ($1 < p < \infty$). Assume A a relatively compact set in $L_p(\Omega)$ with $d(A) > 0$. Then*

$$\begin{aligned}
r(A) &< \frac{1}{\sqrt[p]{2}} \cdot d(A), & 1 < p \leq 2, \quad q := \frac{p}{p-1}; \\
r(A) &< \frac{1}{\sqrt[p]{2}} \cdot d(A), & 2 < p < \infty.
\end{aligned}$$

Theorem 1.5. *Under the assumptions of Theorem 1.1, for every $\varepsilon \in (0, d(A))$, every positive integer m , there exists an m -simplex $\Delta(\varepsilon, m)$ with vertices in A such that each edge of $\Delta(\varepsilon, m)$ has length not less than $d(A) - \varepsilon$.*

Proof. We shall assume A is closed convex, and $r(A) = 1$. From the proof of Theorem 1.1 we derived a sequence $\{y_{s_{n-1}+1}, y_{s_{n-1}+2}, \dots, y_{s_n}\}_{n=2}^\infty$ in A , and a sequence of positive numbers $\{\alpha_{s_{n-1}+1}, \alpha_{s_{n-1}+2}, \dots, \alpha_{s_n}\}_{n=2}^\infty$ (with convention $s_1 = 0$) such that

$$2r_n^\alpha \leq \sum_{i,j \in I_n} \alpha_i \alpha_j \|y_i - y_j\|^\alpha, \quad \sum_{i \in I_n} \alpha_i = 1,$$

where $r_n \in (1 - \frac{1}{n}, 1]$, α as in (1), and $I_n := \{s_{n-1} + 1, s_{n-1} + 2, \dots, s_n\}$.

We denote by

$$\begin{aligned}
T_{nj} &:= \sum_{i \in I_n} \alpha_i \|y_i - y_j\|^\alpha, \\
S_n &:= \{j \in I_n : T_{nj} \geq 2r_n^\alpha (1 - \sqrt{1 - r_n^\alpha})\}, \\
S_n(y_j) &:= \left\{ i \in I_n : \|y_i - y_j\|^\alpha \geq 2 \left(1 - \frac{1}{\sqrt[4]{n}}\right) \right\}, \quad j \in S_n, \\
\hat{S}_n(y_j) &:= \{y_i : i \in S_n(y_j)\}, \quad j \in S_n, \\
\lambda_n &:= \sum_{i \in I_n \setminus S_n} \alpha_i = 1 - \sum_{i \in S_n} \alpha_i.
\end{aligned}$$

Based on the method in proving Theorem 3.4 of [6] one can proceed furthermore as follows. We have

$$\begin{aligned}
2 r_n^\alpha &\leq \sum_{i,j \in I_n} \alpha_i \alpha_j \|y_i - y_j\|^\alpha \\
&= \sum_{j \in S_n} \alpha_j \sum_{i \in I_n} \alpha_i \|y_i - y_j\|^\alpha + \sum_{j \in I_n \setminus S_n} \alpha_j \sum_{i \in I_n} \alpha_i \|y_i - y_j\|^\alpha \\
&\leq 2 \sum_{j \in S_n} \alpha_j + 2 r_n^\alpha (1 - \sqrt{1 - r_n^\alpha}) \sum_{j \in I_n \setminus S_n} \alpha_j \\
&= 2 - 2 \lambda_n (1 - r_n^\alpha + r_n^\alpha \sqrt{1 - r_n^\alpha}) \\
&\leq 2 - 2 \lambda_n \sqrt{1 - r_n^\alpha}.
\end{aligned}$$

Hence $\lambda_n \leq \sqrt{1 - r_n^\alpha} \rightarrow 0$, as $n \rightarrow \infty$. Thus $\lim_{n \rightarrow \infty} \left(\sum_{i \in S_n} \alpha_i \right) = \lim_{n \rightarrow \infty} (1 - \lambda_n) = 1$.

On the other hand

$$2 r_n^\alpha \leq \sum_{i,j \in I_n} \alpha_i \alpha_j \|y_i - y_j\|^\alpha \leq 2 \left(1 - \left(\sum_{i \in I_n} \alpha_i^2 \right) \right) \leq 2 (1 - \alpha_i^2)$$

for every $i \in I_n$. Therefore $\alpha_i \leq \sqrt{1 - r_n^\alpha} \rightarrow 0$ as $n \rightarrow \infty$. One concludes that the cardinality $|S_n|$ of S_n tends to ∞ as $n \rightarrow \infty$. In a similar manner as in the proof of Theorem 3.4 of [6] for every $\varepsilon \in (0, d(A))$ and a given positive integer m we choose n sufficiently large satisfying

$$|S_n| > m, \quad \frac{2 \alpha m}{\sqrt[4]{n}} < 1, \quad 2 \left(1 - \frac{1}{\sqrt[4]{n}} \right) \geq (d(A) - \varepsilon)^\alpha$$

such that for every $1 \leq k \leq m$ and every choice of $i_1, i_2, \dots, i_k \in S_n$ we have

$$\bigcap_{\nu=1}^k \hat{S}_n(y_{i_\nu}) \neq \emptyset.$$

With m and n as above and a fixed $j \in S_n$, setting $z_1 := y_j$, we take consecutively $z_2 \in \hat{S}_n(z_1)$, $z_3 \in \hat{S}_n(z_1) \cap \hat{S}_n(z_2)$, $\dots$, $z_{m+1} \in \bigcap_{k=1}^m \hat{S}_n(z_k)$. One sees that

$$\|z_i - z_j\|^\alpha \geq 2 \left(1 - \frac{1}{\sqrt[4]{n}} \right) \geq (d(A) - \varepsilon)^\alpha.$$

for all $i \neq j$ in $\{1, 2, \dots, m+1\}$, with n sufficiently large. We obtain an m -simplex formed by $z_1, z_2, \dots, z_{m+1}$, whose edges have length not less than $d(A) - \varepsilon$, as claimed.

The proof of Theorem 1.5 is complete.

Remark 1.6. (i) Since for $L_p(\Omega)$ spaces $J_s = J$ the extremal sets in $L_p(\Omega)$ are also self-extremal. Thus we obtain a characterization for extremal sets in $L_p(\Omega)$ via Theorem 1.5 above.

(ii) In particular $\Omega = \mathbb{N}$, $\mu(A) := \text{card}(A)$, $A \subset \mathbb{N}$ leads to the ℓ_p space case (Theorem 3.4 of [6]).

Example 1.7. (i) Let $p \geq 2$, consider a sequence $\{\Omega_n\}_{n=1}^\infty$ consisting of measurable subsets of Ω such that

$$0 < \mu(\Omega_i) < \infty, \quad i = 1, 2, \dots; \quad \Omega_i \cap \Omega_j = \emptyset, \quad \forall i \neq j; \quad \bigcup_{i=1}^{\infty} \Omega_i = \Omega.$$

Let χ_{Ω_i} denote the characteristic function of Ω_i , and set

$$A := \{f_i\}_{i=1}^\infty, \quad \text{where } f_i := \frac{\chi_{\Omega_i}}{\mu(\Omega_i)}.$$

One can check easily that $r(A) = 1$, $d(A) = 2^{\frac{1}{p}}$, hence A is a self-extremal set in $L_p(\Omega)$.

(ii) In the case $1 < p < 2$ we set $B := \{r_i\}_{i=0}^\infty$, where $\{r_i\}_{i=0}^\infty$ is the sequence of Rademacher functions in $L_p[0, 1]$. If $r \in \text{co}\{r_0, r_1, \dots, r_n\}$ and $k \geq n + 1$ then it is easy to see that $d(B) = 2^{1-\frac{1}{p}}$ and

$$\|r - r_k\|_p := \left(\int_0^1 |r - r_k|^p d\mu \right)^{\frac{1}{p}} \geq \left| \int_0^1 (r - r_k) r_k d\mu \right| = 1,$$

hence $r(B) = 1$. Thus B is a self-extremal set in $L_p[0, 1]$ with $1 < p < 2$. This is in contrast to the ℓ_p case ([6]), where we conjectured that there are no (self)-extremal sets in ℓ_p , spaces with $1 < p < 2$.

References

- [1] Amir, D., *On Jung's constant and related constants in normed linear spaces*, Pacific J. Math., **118** (1985), 1–15.
- [2] Dominguez-Benavides, T., *Normal structure coefficients of $L_p(\Omega)$* , Proc. Royal Soc. Edinburgh, **117A** (1990), 299–303.
- [3] Gulevich, N. M., *The radius of a compact set in a Hilbert space*, Zapiski Nauchnykh Seminarov LOMI im. V. A. Steklova, AN SSSR, vol. **164** (1988), 157–158 (Russian).
- [4] Ivanov, V. I., Pichugov, S. A., *Jung's constants of ℓ_p^n -spaces*, Matem. Zametki, **48** (1990), N^o 4, 37–47 (Russian).

- [5] Nguyen-Khac, V., Nguyen-Van, K., *A characterization of extremal sets in Hilbert spaces*, arXiv: math. MG/0203190.
- [6] Nguyen-Khac, V., Nguyen-Van, K., *A geometric characterization of extremal sets in ℓ_p spaces*, arXiv: math. MG/0207304.
- [7] Pichugov, S. A., *Jung's constants of L_p space*, Matem. Zametki, **43** (1988), 604–614 (Russian).
- [8] Pichugov, S. A., *Relative Jung's constants of L_p space*, Ukr. Matem. Zh., **42** (1990), N^o 1, 122–125 (Russian).
- [9] Wells, J. H., Williams, L. R., *Embeddings and Extensions in Analysis*, Springer-Verlag, 1975.